

Application No. 10/009,431

Filed: February 13, 2002

Group Art Unit:1636

Confirmation No.: 2347

AMENDMENTS TO THE CLAIMS

1. (Currently Amended) A pharmaceutical composition comprising a nucleic acid containing a nucleotide sequence encoding the primary amino acid sequence of ~~a protein containing the 7 Cys-knot region~~ GDF-15 protein of the TGF- β superfamily or a functionally active derivative or part thereof having at least a neurotrophic effect on DAergic neurons, wherein the gene coding for the ~~protein containing the 7 Cys-knot region~~ GDF-15 protein of the TGF- β superfamily is transcribed and/or translated in neurons and glial cells, or a vector containing at least the nucleic acid or a protein encoded by the nucleic acid or an antibody or a functional fragment thereof directed against the protein or an antagonist directed to the protein or an agonist as a substitute for the protein, optionally in combination with a pharmaceutically acceptable carrier and/or diluent, for the prevention and/or treatment of neurodegenerative disorders in mammals.

2. (Original) The pharmaceutical composition according to claim 1, wherein the neuron and glial cells are of mammalian origin.

3. (Currently Amended) The pharmaceutical composition according to claim 1, wherein the protein ~~containing the 7 Cys-knot region~~ GDF-15 protein of the TGF- β superfamily protects against neurodegenerative events.

4. (Original) The pharmaceutical composition according to claim 3, wherein the neurodegenerative event is mediated by oxidative

Application No. 10/009,431

Filed: February 13, 2002

Group Art Unit:1636

Confirmation No.: 2347

damage and/or free radical damage and/or mediators and/or executors of neuronal death programs.

5. (Original) The pharmaceutical composition according to claim 4, wherein the mediators of the free radical damage are selected from the group consisting of iron, NO donors, and other free radical donors, and the mediators and executors of neuronal death programs are selected from the group consisting of caspases and pro- and anti-apoptotic members of the bcl-2 family.

6. (Previously Presented) The pharmaceutical composition according to claim 1, wherein the nucleic acid comprises at least the nucleotide sequence shown in Fig. 7A or the nucleotide sequence shown in Fig. 8A or nucleotides 40 to 333 of the nucleotide sequence shown in Fig. 8A or mutants thereof leading to the expression of functionally active polypeptides.

7. (Previously Presented) The pharmaceutical composition according to claim 1, wherein the protein encoded by the nucleic acid comprises at least the primary amino acid sequence shown in Fig. 7B or the primary amino acid sequence shown in Fig. 8B or amino acid residues 14 to 111 of the sequence shown in Fig. 8B as well as homologs thereof having conservative amino acid substitutions.

8. (Previously Presented) The pharmaceutical composition according to claim 1, wherein the mammal is a human.

Application No. 10/009,431
Filed: February 13, 2002
Group Art Unit:1636
Confirmation No.: 2347

9. (Previously Presented) The pharmaceutical composition according to claim 1, wherein the neurodegenerative disorders are selected from the group of acute and/or chronic neurological and psychological disorders.

10. (Original) The pharmaceutical composition of claim 9, wherein the neurological and psychological disorders are caused by stroke, parkinson's disease, Alzheimer's disease or other dementias, infections of the CNS and psychiatric disorders associated with disturbances in CNS transmitter systems.

11. (Original) The pharmaceutical composition according to claim 10, wherein the psychiatric disorders are selected from the group consisting of depression and schizophrenia.

12. (Previously Presented) The pharmaceutical composition according to claim 1, further comprising one or more agents having neurotrophic activity or functionally active derivatives or parts thereof.

13. (Original) The pharmaceutical composition according to claim 12, wherein the agent is a cytokine.

14. (Original) The pharmaceutical composition according to claim 13, wherein the cytokine is selected from the group consisting of GDF, GDNF, TGF, activins, BMP, BDNF, NGF, EGF, CNTF and FGF.

15. (Currently Amended) A diagnostic kit comprising a nucleic acid containing a nucleotide sequence encoding the primary amino acid sequence of a ~~protein containing the 7 Cys knot region~~ GDF-

-4-

WEINCARTEN, SCHURCIN,
GAGNEBIN & LESOVICI LLP
TEL. (617) 542-2290

Application No. 10/009,431

Filed: February 13, 2002

Group Art Unit:1636

Confirmation No.: 2347

15 protein of the TGF- β superfamily or a functionally active derivative or part thereof having at least a neurotrophic effect on DAergic neurons, wherein the gene coding for ~~the protein containing the 7 Cys knot region~~ GDF-15 protein of the TGF- β superfamily is transcribed and/or translated in neurons and glial cells, and/or a vector containing at least the nucleic acid and/or a protein encoded by the nucleic acid and/or an antibody or a functional fragment thereof directed against the protein, for the detection of neurodegenerative disorders in mammals.

16. (Original) The diagnostic kit according to claim 15, wherein the neuron and glial cells are of mammalian origin.

17. (Currently Amended) The diagnostic kit according to claim 15, wherein the ~~protein containing the 7 Cys knot region~~ GDF-15 protein of the TGF- β superfamily protects against neurodegenerative events.

18. (Original) The diagnostic kit according to claim 17, wherein the neurodegenerative event is mediated by oxidative damage and/or free radical damage and/or mediators and/or executors of neuronal death programs.

19. (Original) The diagnostic kit according to claim 18, wherein the mediators of the free radical damage are selected from the group consisting of iron, NO donors, and other free radical donors, and the mediators and executors of neuronal death programs are selected from the group consisting of caspases and pro- and anti-apoptotic members of the bcl-2 family.

-5-

WEINSTEIN, SCHUBIN,
CAGNINI & LEBOVITZ LLP
TEL. (617) 542 2290
FAX (617) 542 2290

Application No. 10/009,431

Filed: February 13, 2002

Group Art Unit:1636

Confirmation No.: 2347

20. (Previously Presented) The diagnostic kit according to claim 15, wherein the nucleic acid comprises at least the nucleotide sequence shown in Fig. 7A or the nucleotide sequence shown in Fig. 8A or nucleotides 40 to 333 of the nucleotide sequence shown in Fig. 8A or mutants thereof leading to the expression of functionally active polypeptides.

21. (Previously Presented) The diagnostic kit according to claim 15, wherein the protein encoded by the nucleic acid comprises at least the primary amino acid sequence shown in Fig. 7B or the primary amino acid sequence shown in Fig. 8B or amino acid residues 14 to 111 of the sequence shown in Fig. 8B as well as homologs thereof having conservative amino acid substitutions.

22. (Previously Presented) The diagnostic kit according to claim 19, wherein the mammal is a human.

23. (Currently Amended) The pharmaceutical composition according to claim 2, wherein ~~the protein containing the 7 Cys knot region~~ GDF-15 protein of the TGF- β superfamily protects against neurodegenerative events.

24. (Currently Amended) The diagnostic kit according to claim 16, wherein ~~the protein containing the 7 Cys knot region~~ GDF-15 protein of the TGF- β superfamily protects against neurodegenerative events.

25. (Currently Amended) The pharmaceutical composition according to claim 2, wherein:

-6-

WEINGARTEN, SCHURGIN,
CAGNESIN & LEBOVICI LLP
TEL. (517) 542-2390
FAX (517) 542-2390

Application No. 10/009,431
Filed: February 13, 2002
Group Art Unit:1636
Confirmation No.: 2347

~~the protein containing the 7-Cys knot region~~ GDF-15 protein of
the TGF- β

superfamily protects against neurodegenerative events;
the neurodegenerative event is mediated by oxidative damage
and/or free radical damage and/or mediators and/or executors of
neuronal death programs;

the mediators of the free radical damage are selected from
the group consisting of iron, NO donors, and other free radical
donors, and the mediators and executors of neuronal death
programs are selected from the group consisting of caspases and
pro- and anti-apoptotic members of the bcl-2 family;
either the nucleic acid comprises at least the nucleotide
sequence shown in Fig. 7A or the nucleotide sequence shown in
Fig. 8A or nucleotides 40 to 333 of the nucleotide sequence
shown in Fig. 8A or mutants thereof leading to the expression of
functionally active polypeptides; or the protein encoded by the
nucleic acid comprises at least the primary amino acid sequence
shown in Fig. 7B or the primary amino acid sequence shown in
Fig. 8B or amino acid residues 14 to 111 of the sequence shown
in Fig. 8B as well as homologs thereof having conservative amino
acid substitutions;

the mammal is a human;

the neurodegenerative disorders are selected from the group
of acute and/or chronic neurological and psychological
disorders;

the neurological and psychological disorders are caused by
stroke, parkinson's disease, Alzheimer's disease or other
dementias, infections of the CNS and psychiatric disorders
associated with disturbances in CNS transmitter systems;

Application No. 10/009,431

Filed: February 13, 2002

Group Art Unit:1636

Confirmation No.: 2347

the psychiatric disorders are selected from the group consisting of depression and schizophrenia;

the agent is a cytokine selected from the group consisting of GDF, GDNF, TGF, activins, BMP, BDNF, NGF, EGF, CNTF and FGF; and

further comprising one or more agents having neurotrophic activity or functionally active derivatives or parts thereof.

26. (Currently Amended) The diagnostic kit according to claim 16, wherein:

~~the protein containing the 7 Cys knot region~~ GDF-15 protein of the TGF- β superfamily protects against neurodegenerative events;

the neurodegenerative event is mediated by oxidative damage and/or free radical damage and/or mediators and/or executors of neuronal death programs;

the mediators of the free radical damage are selected from the group consisting of iron, NO donors, and other free radical donors, and the mediators and executors of neuronal death programs are selected from the group consisting of caspases and pro- and anti-apoptotic members of the bcl-2 family;

wherein the nucleic acid comprises at least the nucleotide sequence shown in Fig. 7A or the nucleotide sequence shown in Fig. 8A or nucleotides 40 to 333 of the nucleotide sequence shown in Fig. 8A or mutants thereof leading to the expression of functionally active polypeptides;

the protein encoded by the nucleic acid comprises at least the primary amino acid sequence shown in Fig. 7B or the primary amino acid sequence shown in Fig. 8B or amino acid residues 14 to 111 of the sequence shown in Fig. 8B as well as homologs thereof having conservative amino acid substitutions; and

-8-

Application No. 10/009,431
Filed: February 13, 2002
Group Art Unit:1636
Confirmation No.: 2347

the mammal is a human.